



Associate : PKM

Sample Type : O.S.S.

Name : MAYA DALUI

Age / Gender : 50Y / FEMALE

Received On : 22/03/2026

ID Number : LSD//64612/10

UHID : 3260322064612 Reported On : 22/03/2026

Referred By : DR K MD YEZDANI

User:isd
ClinicPro / LifeSafe
Page 1 of 1

DEPARTMENT OF CLINICAL BIOCHEMISTRY

<u>Test Description</u>	<u>Result</u>	<u>Unit</u>	<u>Bio. Ref. Interval</u>
<u>LFT (Liver Function Test)</u>			
Bilirubin Total <i>Methodology : Jendrassik Spectrophotometry</i> <i>Specimen : Serum</i>	0.81	mg/dl	Adult: Upto 1.2 Infant: 0.2 - 8.0
Bilirubin Direct <i>Methodology : DIAZO</i>	0.24	mg/dl	0.0 - 0.4
Bilirubin Indirect	0.57	mg/dl	upto 0.75
Total Protein <i>Methodology : BIURET</i> <i>Specimen : Serum</i>	6.19	gm/dl	6.0-8.0
Albumin <i>Methodology : Bromocresol Green</i> <i>Specimen : Serum</i>	3.90	gm/dl	3.5 - 5.2
Globulin <i>Specimen : Serum</i>	2.29	gm/dl	2.3 - 3.6
A G Ratio	1.70		1.0 - 2.3
ALP (Alkaline Phosphatase) <i>Methodology : IFCC</i>	177.02	U/L	Adults : 100 - 290 U/L Children: 180 - 1200 U/L
SGOT / AST <i>Methodology : IFCC</i> <i>Specimen : Serum</i>	43.0	IU/L	upto 46
SGPT / ALT <i>Methodology : IFCC</i> <i>Specimen : Serum</i>	29.1	U/L	upto 49

Comment:

AST (SGOT) and ALT (SGPT) are sensitive indicators of liver damage from different types of disease. The highest levels of AST and ALT are found with disorders that cause the death of numerous liver cells (extensive hepatic necrosis). There are several drugs that may cause abnormal liver enzymes levels.

End of Report

